

### ***Disclosure of the Invention***

### Background of the Invention

The results of investigations directed to the understanding of pathogenesis of mental disorders have shown that a disorder in the serotonin equilibrium plays an important role in various diseases. The monoamine-deficiency hypothesis was one of the first explanations, wherein the symptoms of depression were connected to a reduction in the neurotransmission of monoamines, especially serotonin (5-HT) and noradrenaline, which was also confirmed by neurochemical tests as well as by a successful treatment of the patients with substances increasing monoaminergic neurotransmission (*Expert Opin. Investig. Drugs* **2003**, 12, 531–543). In addition to the serotonergic and noradrenergic

For treatment of pathological CNS disorders and particularly for mental disorders, the most frequently applied medicines ~~aFor treatment of pathological CNS disorders and particularly in the therapy of mental disorders a significant role as the most frequently applied medicines is given to substances that, according to their structure, are polycyclic compounds (benzodiazepines, tricyclic and tetracyclic antidepressants, monoamino oxidase (MAO) inhibitors, selective inhibitors of serotonin reabsorption etc.).~~

{W:\03818\0204416us0\00747961.DOC }

However, the known medicines for CNS disorders and particularly in the therapy of mental disorders are associated with a wide range of adverse effects. There is thus, there a need for a safe and effective treatment of diseases and disorders of CNS.

Moreover, no compound representing the subject matter of the present invention has been described as effective in the treatment of diseases and disorders of CNS. Consequently, the use of 1-aza-2-oxa-dibenzo[*e,h*]azulenes and of their pharmaceutically acceptable salts and solvates for the manufacture ~~or use in a pharmaceutical compositions~~ for the treatment and prevention of diseases, damages and


disorders of the central nervous system caused by disorders of neurochemical equilibrium has hitherto been neither disclosed nor suggested.

**Summary of the Invention**

The compounds from the class of 1-aza-2-oxa-dibenzo[*e,h*]azulenes represented by the formula **I**, differ structurally from the art-known tetracyclic compounds acting upon CNS by an unsaturated tetracyclic structure since they contain an isoxazole ring as the fourth ring, whereas the art-known tetracyclic compounds acting upon CNS (WO 99/19317, WO 97/38991; Sperling, W.; Demling, J. *Drugs Today* **1997**, 33, 95–102) contain at least one saturated ring in their structure, and are further distinguished by valuable pharmacological and physicochemical properties.

The compounds represented by the formula **I**, which are the subject matter of the present invention, isomeric forms of such compounds, their pharmaceutically acceptable salts and solvates and pharmaceutical composition comprising them are not believed to have been previously described. Moreover, no compound representing the subject matter of the present invention has been described as effective in the treatment of diseases and disorders of CNS.

The present invention relates to the compounds from the class of 1-aza-2-oxa-dibenzo[*e,h*]azulenes of the general formula **I**:



wherein

X means CH<sub>2</sub> or a heteroatom selected from the group consisting of O, S, S(=O), S(=O)<sub>2</sub> and NR<sup>a</sup>, wherein R<sup>a</sup> is hydrogen or a substituent selected from the group consisting of C<sub>1</sub>-C<sub>3</sub>-alkyl, C<sub>1</sub>-C<sub>3</sub>-alkanoyl, C<sub>1</sub>-C<sub>7</sub>-alkoxycarbonyl, C<sub>7</sub>-C<sub>10</sub>-arylalkyloxycarbonyl, C<sub>7</sub>-C<sub>10</sub>-aroyl, C<sub>7</sub>-C<sub>10</sub>-arylalkyl, C<sub>3</sub>-C<sub>7</sub>-alkylsilyl and C<sub>5</sub>-C<sub>10</sub>-alkylsilylalkyloxyalkyl;


{W:\03818\0204416us0\00747961.DOC {TEXT MARKED WITH MICROSOFT WORD}} {TEXT MARKED WITH MICROSOFT WORD}}

Y and Z independently from each other mean one or more identical or different substituents linked to any available carbon atom selected from the group consisting of hydrogen, halogen, C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>2</sub>-C<sub>4</sub>-alkenyl, C<sub>2</sub>-C<sub>4</sub>-alkynyl, halo-C<sub>1</sub>-C<sub>4</sub>-alkyl, hydroxy, C<sub>1</sub>-C<sub>4</sub>-alkoxy, trifluoromethoxy, C<sub>1</sub>-C<sub>4</sub>-alkanoyl, amino, amino-C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>1</sub>-C<sub>4</sub>-alkylamino, *N*-(C<sub>1</sub>-C<sub>4</sub>-alkyl)amino, *N,N*-di(C<sub>1</sub>-C<sub>4</sub>-alkyl)amino, thiol, C<sub>1</sub>-C<sub>4</sub>-alkylthio, sulfonyl, C<sub>1</sub>-C<sub>4</sub>-alkylsulfonyl, sulfinyl, C<sub>1</sub>-C<sub>4</sub>-alkylsulfinyl, carboxy, C<sub>1</sub>-C<sub>4</sub>-alkoxycarbonyl, cyano and nitro;

R<sup>1</sup> means hydrogen, halogen, C<sub>1</sub>-C<sub>7</sub>-alkyl optionally substituted with one, two, three or more substituents selected from the group consisting of halogen atom, hydroxy, C<sub>1</sub>-C<sub>4</sub> alkoxy, thiol, C<sub>1</sub>-C<sub>4</sub> alkylthio, amino, *N*-(C<sub>1</sub>-C<sub>4</sub>) alkylamino, *N,N*-di(C<sub>1</sub>-C<sub>4</sub>-alkyl)-amino, sulfonyl, C<sub>1</sub>-C<sub>4</sub> alkylsulfonyl, sulfinyl and C<sub>1</sub>-C<sub>4</sub> alkylsulfinyl; C<sub>2</sub>-C<sub>7</sub>-alkenyl optionally substituted with one, two, three or more halogen atoms; C<sub>2</sub>-C<sub>7</sub>-alkynyl; monocyclic or bicyclic aryl group having from 6 to 10 carbon atoms and altering double bond and said group can be optionally substituted with one or two substituents selected from the group consisting of fluoro, chloro, C<sub>1</sub>-C<sub>4</sub> alkyl, cyano, nitro, hydroxy, C<sub>1</sub>-C<sub>4</sub> alkoxy, thiol, C<sub>1</sub>-C<sub>4</sub> alkylthio, amino, *N*-(C<sub>1</sub>-C<sub>4</sub>) alkylamino, *N,N*-di(C<sub>1</sub>-C<sub>4</sub>-alkyl)-amino, sulfonyl, C<sub>1</sub>-C<sub>4</sub> alkylsulfonyl, sulfinyl, C<sub>1</sub>-C<sub>4</sub> alkylsulfinyl and can be linked to the rest of the molecule by any available carbon atom via direct bond or via C<sub>1</sub>-C<sub>4</sub> alkylene group; monocyclic or bicyclic heteroaryl having the meaning of aromatic and partially aromatic groups of a monocyclic or bicyclic ring with 4 to 12 carbon atoms and at least one of them being heteroatom selected from the group consisting of O, S and N wherein available carbon or nitrogen represent the binding site of the group to the rest of the molecule either via direct bond or via C<sub>1</sub>-C<sub>4</sub> alkylene group and where said heteroaryl can be optionally substituted with fluoro, chloro, C<sub>1</sub>-C<sub>4</sub> alkyl, cyano, nitro, hydroxy, C<sub>1</sub>-C<sub>4</sub> alkoxy, thiol, C<sub>1</sub>-C<sub>4</sub> alkylthio, amino, *N*-(C<sub>1</sub>-C<sub>4</sub>) alkylamino, *N,N*-di(C<sub>1</sub>-C<sub>4</sub>-alkyl)-amino, sulfonyl, C<sub>1</sub>-C<sub>4</sub> alkylsulfonyl, sulfinyl, C<sub>1</sub>-C<sub>4</sub> alkylsulfinyl; five-member or six-member fully saturated or partly unsaturated heterocycle group containing at least one hetero atom selected from the group consisting of O, S and N wherein available carbon or nitrogen represent the binding site of the group to the rest of the molecule either via direct bond or via C<sub>1</sub>-C<sub>4</sub> alkylene group and where said heterocycle can be optionally substituted with fluoro, chloro, C<sub>1</sub>-C<sub>4</sub> alkyl, cyano, nitro, hydroxy, C<sub>1</sub>-C<sub>4</sub> alkoxy, thiol, C<sub>1</sub>-C<sub>4</sub> alkylthio, amino, *N*-(C<sub>1</sub>-C<sub>4</sub>) alkylamino, *N,N*-di(C<sub>1</sub>-C<sub>4</sub>-alkyl)-amino, sulfonyl, C<sub>1</sub>-C<sub>4</sub> alkylsulfonyl, sulfinyl, C<sub>1</sub>-C<sub>4</sub> alkylsulfinyl; hydroxy; hydroxy-C<sub>2</sub>-C<sub>7</sub>-alkenyl; hydroxy-C<sub>2</sub>-C<sub>7</sub>-alkynyl; C<sub>1</sub>-C<sub>7</sub>-alkoxy; thiol; thio-C<sub>2</sub>-C<sub>7</sub>-alkenyl; thio-C<sub>2</sub>-C<sub>7</sub>-alkynyl; C<sub>1</sub>-C<sub>7</sub>-alkylthio; amino; *N*-(C<sub>1</sub>-C<sub>7</sub>-alkyl)amino; *N,N*-di(C<sub>1</sub>-C<sub>7</sub>-alkyl)amino; C<sub>1</sub>-C<sub>7</sub>-alkylamino; amino-C<sub>2</sub>-C<sub>7</sub>-alkenyl; amino-C<sub>2</sub>-C<sub>7</sub>-alkynyl; amino-C<sub>1</sub>-C<sub>7</sub>-alkoxy; C<sub>1</sub>-C<sub>7</sub>-alkanoyl; C<sub>7</sub>-C<sub>10</sub>-aroyl; oxo-C<sub>1</sub>-C<sub>7</sub>-alkyl; C<sub>1</sub>-C<sub>7</sub>-alkanoyloxy; carboxy; an optionally substituted C<sub>1</sub>-C<sub>7</sub>-alkyloxycarbonyl; an optionally substituted C<sub>7</sub>-C<sub>10</sub>-aryloxycarbonyl; carbamoyl; *N*-(C<sub>1</sub>-C<sub>7</sub>-alkyl)carbamoyl; *N,N*-di(C<sub>1</sub>-C<sub>7</sub>-

alkyl)carbamoyl; cyano; cyano-C<sub>1</sub>-C<sub>7</sub>-alkyl; sulfonyl; C<sub>1</sub>-C<sub>7</sub>-alkylsulfonyl; sulfinyl; C<sub>1</sub>-C<sub>7</sub>-alkylsulfinyl; nitro;

or a substituent represented with the formula II:



## II

wherein

R<sup>2</sup> and R<sup>3</sup> simultaneously or independently from each other ~~have the meaning of~~ are hydrogen, C<sub>1</sub>-C<sub>4</sub>-alkyl, aryl having the meaning as defined above; or together with N ~~have the meaning of~~ are optionally substituted heterocycle or heteroaryl wherein heterocycle relates to five-membered or six-membered fully saturated or partly unsaturated heterocycle group containing at least one hetero atom selected from the group consisting of O, S and N and where said heterocycle can be optionally substituted with one or two substituents which are selected from halogen, C<sub>1</sub>-C<sub>4</sub> alkyl, cyano, nitro, hydroxy, C<sub>1</sub>-C<sub>4</sub> alkoxy, thiol, C<sub>1</sub>-C<sub>4</sub> alkylthio, amino, *N*-(C<sub>1</sub>-C<sub>4</sub>) alkylamino, *N,N*-di(C<sub>1</sub>-C<sub>4</sub>-alkyl)-amino, sulfonyl, C<sub>1</sub>-C<sub>4</sub> alkylsulfonyl, sulfinyl, C<sub>1</sub>-C<sub>4</sub> alkylsulfinyl and heteroaryl relates to aromatic and partially aromatic groups of a monocyclic or bicyclic ring with 4 to 12 carbon atoms and at least one of them being heteroatom selected from the group consisting of O, S and N and where said heteroaryl can be optionally substituted with one or two substituents which are selected from halogen, C<sub>1</sub>-C<sub>4</sub> alkyl, cyano, nitro, hydroxy, C<sub>1</sub>-C<sub>4</sub> alkoxy, thiol, C<sub>1</sub>-C<sub>4</sub> alkylthio, amino, *N*-(C<sub>1</sub>-C<sub>4</sub>) alkylamino, *N,N*-di(C<sub>1</sub>-C<sub>4</sub>-alkyl)-amino, sulfonyl, C<sub>1</sub>-C<sub>4</sub> alkylsulfonyl, sulfinyl, C<sub>1</sub>-C<sub>4</sub> alkylsulfinyl;

m ~~has the meaning of~~ is an integer from 1 to 3;

Q ~~has the meaning of~~ is oxygen, sulfur or nitrogen;

and to their pharmaceutically acceptable salts and solvates, as well as to pharmaceutical compositions containing one or more of the foregoing compounds in an amount effective to treat and prevent diseases, damages and disorders of the central nervous system caused by disorders of neurochemical equilibrium of biogenic amines or other neurotransmitters.

When X ~~has the meaning of~~ is NR<sup>a</sup>, R<sup>a</sup> relates to hydrogen or group selected from the C<sub>1</sub>-C<sub>3</sub>-alkyl (preferably methyl or ethyl), C<sub>1</sub>-C<sub>3</sub>-alkanoyl (preferably formyl or acetyl), C<sub>1</sub>-C<sub>7</sub>-alkoxycarbonyl

{W:\03818\0204416us0\00747961.DOC /AMT/03/03/2004/00747961.DOC}

When R<sup>2</sup> and R<sup>3</sup> together with N have the meaning of ~~are~~ heteroaryl or heterocycle, this means that such heteroaryls or heterocycles have at least one carbon atom replaced by a nitrogen atom through which the groups are linked to the rest of the molecule. Examples of such groups are morpholin-4-yl, piperidin-1-yl, pyrrolidin-1-yl, imidazol-1-yl or piperazin-1-yl.

In one embodiment of the present invention preferred compounds of formula **I** are those wherein X represents O or S.

In another embodiment of the present invention preferred compounds of formula **I** are those wherein Y and Z independently from each other mean one or more identical or different substituents linked to any available carbon atom selected from the group consisting of hydrogen, fluorine, chlorine, bromine, C<sub>1</sub>-C<sub>4</sub>-alkyl (preferably methyl, ethyl, propyl or isopropyl), halo-C<sub>1</sub>-C<sub>4</sub>-alkyl (preferably trifluoromethyl), hydroxy, C<sub>1</sub>-C<sub>4</sub>-alkoxy (preferably methoxy), trifluoromethoxy, C<sub>1</sub>-C<sub>4</sub>-alkanoyl (preferably formyl or acetyl), amino, amino-C<sub>1</sub>-C<sub>4</sub>-alkyl (preferably aminomethyl), *N*-(C<sub>1</sub>-C<sub>4</sub>-alkyl)amino (preferably *N*-methyl or *N*-ethyl), *N,N*-di(C<sub>1</sub>-C<sub>4</sub>-alkyl)amino (preferably dimethylamino or diethylamino), thiol, C<sub>1</sub>-C<sub>4</sub>-alkylthio (preferably methylthio), cyano and nitro.

In yet another embodiment of the present invention preferred compounds of formula **I** are those wherein R<sup>1</sup> has the meaning of hydrogen, halogen, C<sub>1</sub>-C<sub>7</sub>-alkyl optionally substituted with one, two, three or more substituents selected from the group consisting of halogen atom, hydroxy, C<sub>1</sub>-C<sub>4</sub> alkoxy, thiol, C<sub>1</sub>-C<sub>4</sub> alkylthio, amino, *N*-(C<sub>1</sub>-C<sub>4</sub>) alkylamino and *N,N*-di(C<sub>1</sub>-C<sub>4</sub>-alkyl)-amino; monocyclic or bicyclic aryl group having from 6 to 10 carbon atoms and altering double bond and said group can be optionally substituted with one or two substituents selected from the group consisting of fluoro, chloro, C<sub>1</sub>-C<sub>4</sub> alkyl, cyano, nitro, hydroxy, C<sub>1</sub>-C<sub>4</sub> alkoxy, thiol, C<sub>1</sub>-C<sub>4</sub> alkylthio, amino, *N*-(C<sub>1</sub>-C<sub>4</sub>) alkylamino and *N,N*-di(C<sub>1</sub>-C<sub>4</sub>-alkyl)-amino and can be linked to the rest of the molecule by any available carbon atom via direct bond or via C<sub>1</sub>-C<sub>4</sub> alkylene group; monocyclic or bicyclic heteroaryl having the meaning of aromatic and partially aromatic groups of a monocyclic or bicyclic ring with 4 to 12 carbon atoms and at least one of them being heteroatom selected from the group consisting of O, S and N wherein available carbon or nitrogen represent the binding site of the group to the rest of the molecule either via direct bond or via C<sub>1</sub>-C<sub>4</sub> alkylene group and where said heteroaryl can be optionally substituted with fluoro, chloro, C<sub>1</sub>-C<sub>4</sub> alkyl, cyano, nitro, hydroxy, C<sub>1</sub>-C<sub>4</sub>

$$Q-(CH_2)_m-N\begin{matrix} R^2 \\ R^3 \end{matrix}$$

wherein

m ~~has the meaning of~~ is an integer from 1 to 3;

☐ ~~has the meaning of~~ is oxygen.

3-methyl-2-oxa-8-thia-1-aza-dibenzo[*e,h*]azulene;  
11-chloro-3-methyl-2-oxa-8-thia-1-aza-dibenzo[*e,h*]azulene;  
3-methyl-2,8-dioxa-1-aza-dibenzo[*e,h*]azulene;  
3-bromomethyl-2-oxa-8-thia-1-aza-dibenzo[*e,h*]azulene;  
3-bromomethyl-11-chloro-2-oxa-8-thia-1-aza-dibenzo[*e,h*]azulene;  
3-bromomethyl-2,8-dioxa-1-aza-dibenzo[*e,h*]azulene;  
dimethyl-[2-(2-oxa-8-thia-1-aza-dibenzo[*e,h*]azulen-3-ylmethoxy)-ethyl]-amine;  
dimethyl-[3-(2-oxa-8-thia-1-aza-dibenzo[*e,h*]azulen-3-ylmethoxy)-propyl]-amine;



The term "halo", "hal" or "halogen" relates to a halogen atom which may be fluorine, chlorine, bromine or iodine.

The term "haloalkyl" relates to alkyl groups which must be substituted with at least one halogen atom. The most frequent haloalkyls are e.g. chloromethyl, dichloromethyl, trifluoromethyl or 1,2-dichloropropyl.

The term "alkynylalkynyl" relates to alkynylalkynyl groups having the meaning of hydrocarbon radicals, which are straight or branched and contain at least one and at most two carbon-carbon triple bonds. The most frequent alkynylalkynyls are e.g. ethynyl, propynyl or butynyl.

The term "aryl" relates to groups having the meaning of an aromatic ring, e.g. phenyl, as well as to fused aromatic rings. Aryl contains one ring with at least 6 carbon atoms or two rings with totally total of 10 carbon atoms and with alternating double (resonant) bonds between carbon atoms. The most frequently used aryls are e.g. phenyl or naphthyl. In general, aryl groups may be linked to the rest

The term "heterocycle" relates to five-member or six-member, completely saturated or partly unsaturated heterocyclic groups containing at least one hetero atom such as O, S or N, and the available nitrogen atom or carbon atom is the binding site of the group to the rest of the molecule either via a direct bond or via a C<sub>1</sub>-C<sub>4</sub>-alkylene group defined earlier. The most frequent examples are morpholinyl, piperidyl, piperazinyl, pyrrolidinyl, pirazinyl or imidazolyl.

The term "aroyl" group relates to aromatic acyl groups such as benzoyl.

The term "optionally substituted alkenyl" relates to alkenyl groups optionally additionally substituted with one, two or three halogen atoms. Such substituents may be e.g. 2-chloroethenyl, 1,2-dichloroethenyl or 2-bromo-propen-1-yl.

The term "optionally substituted aryl, heteroaryl or heterocycle" relates to aryl, heteroaryl or heterocyclic groups which may be optionally additionally substituted with one or two substituents. The substituents may be halogen (preferably chlorine or fluorine), C<sub>1</sub>-C<sub>4</sub>-alkyl (preferably methyl, ethyl or isopropyl), cyano, nitro, hydroxy, C<sub>1</sub>-C<sub>4</sub>-alkoxy (preferably methoxy or ethoxy), thiol, C<sub>1</sub>-C<sub>4</sub>-

The present invention also relates to all possible tautomeric forms of particular compounds of the formula I.

The term “salts” can include acid addition salts or addition salts of free bases. Examples of acids which may be employed to form pharmaceutically acceptable acid addition salts include but are not limited to salts derived from nontoxic inorganic acids such as nitric, phosphoric, sulfuric, or hydrobromic, hydroiodic, hydrofluoric, phosphorous, as well as salts derived from nontoxic organic acids such as aliphatic mono- and dicarboxylic acids, phenyl-substituted alkanolic acids, hydroxyl alkanolic acids, alkanedioic acids, aromatic acids, aliphatic and aromatic sulfonic acids, and acetic, maleic, succinic, or citric acids. Non-limiting examples of such salts include napadisylate, besylate, sulfate, pyrosulfate, bisulfate, sulfite, bisulfite, nitrate, phosphate, monohydrogenphosphate, dihydrogenphosphate, metaphosphate, pyrophosphate, chloride, bromide, iodide, acetate, trifluoroacetate, propionate, caprylate, isobutyrate, oxalate, malonate, succinate, suberate, sebacate, fumarate, maleate, mandelate, benzoate, chlorobenzoate, methylbenzoate, dinitrobenzoate, phthalate, benzenesulfonate, toluenesulfonate, phenylacetate, citrate, lactate, maleate, tartrate, methanesulfonate, and the like. Also contemplated are salts of amino acids such as arginate and the like and gluconate, galacturonate (see, for example, Berge S. M. et al. “Pharmaceutical Salts,” J. of Pharma. Sci., 1977; 66:1).

[illegible]

physical properties such as solubility in polar solvents, but otherwise the salts are equivalent to their respective free base for purposes of the present invention.

Pharmaceutically acceptable base addition salts are formed with metals or amines, such as alkali and alkaline earth metals or organic amines. Examples of metals used as cations are sodium, potassium, magnesium, calcium, and the like. Examples of suitable amines are N,N'-dibenzylethylenediamine, chloroprocaine, choline, diethanolamine, dicyclohexylamine, ethylenediamine, N-methylglucamine, and procaine.

The base addition salts of said acidic compounds are prepared by contacting the free acid form with a sufficient amount of the desired base to produce the salt in the conventional manner. The free acid form may be regenerated by contacting the salt form with an acid and isolating the free acid.


Preferred pharmaceutically acceptable salts according to invention relate to salts of the formula **I** and include e.g. salts with C<sub>1</sub>-C<sub>4</sub>-alkylhalides (preferably methyl bromide, methyl chloride) (quaternary ammonium salts), with inorganic acids (hydrochloric, hydrobromic, phosphoric, metaphosphoric, nitric or sulfuric acids) or with organic acids (tartaric, acetic, citric, maleic, lactic, fumaric, benzoic, succinic, methane sulfonic or *p*-toluene sulfonic acids).

Pharmaceutically acceptable solvates formed by the compounds represented by formula **I** or their salts relate to hydrates, ethanolates and similar (most frequently hydrates).

The phrase "pharmaceutically acceptable", as used in connection with compositions of the invention, refers to molecular entities and other ingredients of such compositions that are physiologically tolerable and do not typically produce untoward reactions when administered to a mammal (e.g., human). Preferably, as used herein, the term "pharmaceutically acceptable" means approved by a regulatory agency of the Federal or a state government or listed in the U.S. Pharmacopoeia or other generally recognized pharmacopoeias for use in mammals, and more particularly in humans.

A further object of the present invention relates to the preparation of the compounds of the formula **I** according to the following processes:

a) condensation of compound **Ia**:




1

of th

---

Scheme I



{W:\03818\0204416us0\00747961.DOC {UNCLASSIFIED//NOFORN}} UNCLASSIFIED//NOFORN

The starting alcohols, thioalcohols or the compounds of the formula **IIa** are commercially available substances or are prepared according to methods disclosed for the preparation of analogous compounds.

The starting compounds, alcohols of the formula **IIb**, may be obtained by the action of water, ammonia or hydrogen sulfide upon halides of formula **Ia** in a manner disclosed in the literature. The starting optionally selected halides or compounds of the formula **IIb** are already known or are prepared according to methods disclosed for the preparation of analogous compounds.

A further general example of transformation is formylation of the compounds of the formula **I** by processes such as e.g. Vilsmeier acylation or reaction of *n*-BuLi and dimethylformamide. The reaction conditions of these processes are known in the literature.

{W:\03818\0204416us0\00747961.DOC 4/20/2004 10:00:00 AM}

By a selective oxidation of alkylthio group, alkylsulfinyl or alkylsulfonyl groups may be prepared.

By the reduction of the compounds with a nitro group, the preparation of amino compounds is made possible. The reaction is carried out under usual conditions of catalytic hydrogenation or electrochemically. By catalytic hydrogenation using palladium on carbon, alkenyl substituents may be converted to alkyl ones or nitrile group can be converted to aminoalkyl.

Various substituents of the aromatic structure in the compounds of the formula **I** may be introduced by standard substitution reactions or by usual changes of individual functional groups. Examples of such reactions are aromatic substitutions, alkylations, halogenation, hydroxylation as well as oxidation or reduction of substituents. Reagents and reaction conditions are known from the literature. Thus e.g. by aromatic substitution a nitro group is introduced in the presence of concentrated nitric acid and sulfuric acid. By using acyl halides or alkyl halides, the introduction of an acyl group or an alkyl group is made possible. The reaction is carried out in the presence of Lewis acids such as aluminum- or iron-trichloride in conditions of Friedel-Craft reaction. By the reduction of the nitro group, an amino group is obtained, which is by a diazotizing reaction converted to a suitable starting group, which may be replaced with one of the following groups: H, CN, OH, Hal.

In order to prevent undesired interaction in chemical reactions, it is often necessary to protect certain groups such as e.g. hydroxy, amino, thio or carboxy. For this purpose a great number of protecting groups may be used [Green TW, Wuts PGH, *Protective Groups in Organic Synthesis*, John Wiley and Sons, 1999]) and the choice, use and elimination thereof are conventional methods in chemical synthesis.

[illegible]



A convenient protection for amino or alkylamino groups are groups such as e.g. alkanoyl (acetyl), alkoxy carbonyl (methoxy carbonyl, ethoxy carbonyl or *tert*-butoxy carbonyl); arylmethoxy carbonyl (benzyloxy carbonyl), aroyl (benzoyl) or alkylsilyl (trimethylsilyl or trimethylsilylethoxymethyl) groups. The conditions of removing a protecting group depend upon the choice and the characteristics of this group. Thus e.g. acyl groups such as alkanoyl, alkoxy carbonyl or aroyl may be eliminated by hydrolysis in the presence of a base (sodium hydroxide or potassium hydroxide), *tert*-butoxy carbonyl or alkylsilyl (trimethylsilyl) may be eliminated by treatment with a suitable acid (hydrochloric, sulfuric, phosphoric or trifluoroacetic acid), whereas arylmethoxy carbonyl group (benzyloxy carbonyl) may be eliminated by hydrogenation using a catalyst such as palladium on carbon.

The compounds of the present invention are especially effective in treating those diseases and disorders where the neurochemical equilibrium of biogenic amines such as serotonin, norepinephrine and dopamine was disturbed and which may be caused by unbalanced (too big or too small) synthesis, irregularities in storing, releasing, metabolizing and/or reabsorption of a certain neurotransmitter.

It has been found that the compounds of the present invention exhibit a significant binding affinity and have a high degree of selectivity to serotonin receptors, especially to 5-HT<sub>2A</sub> and 5-HT<sub>2C</sub>, as well as for the  $\sigma_1$  receptor.

In one embodiment of the present invention the compound of formula **I**, or salt, or solvate thereof show binding affinity to 5-HT<sub>2A</sub> and 5-HT<sub>2C</sub> serotonin receptors in the concentration expressed as an IC<sub>50</sub> value less than 1  $\mu$ M and having K<sub>i</sub> value less than 1  $\mu$ M.

In another embodiment of the present invention the compound of formula I, or salt, or solvate thereof show binding affinity to 5-HT<sub>2A</sub> serotonin receptor in the concentration expressed as an IC<sub>50</sub> value less than about 200 nM and having K<sub>i</sub> value less than about 100 nM.

In yet another embodiment of the present invention the compound of formula **I**, or salt, or solvate thereof show binding affinity to 5-HT<sub>2C</sub> serotonin receptor in the concentration expressed as an IC<sub>50</sub> value less than about 200 nM and having K<sub>i</sub> value less than about 100 nM.

It has been found that the compounds of the present invention exhibit a significant binding affinity to the  $\sigma_{1\alpha}$  receptor.

In one embodiment of the present invention the compound of formula I, or salt, or solvate thereof show binding affinity to ~~the  $\sigma_{1\sigma 1}$~~  receptor in the concentration expressed as an  $IC_{50}$  value less than 1  $\mu M$  and having  $K_i$  value less than 1  $\mu M$ .

In another embodiment of the present invention the compound of formula I, or salt, or solvate thereof show binding affinity to ~~the  $\sigma_{1\sigma 1}$~~  receptor in the concentration expressed as an  $IC_{50}$  value less than about 200 nM and having  $K_i$  value less than about 100 nM.

Since serotonin receptors are crucial in pathophysiology of a series of CNS disorders (directly or indirectly by participating in the activation of some other neurotransmitter e.g. dopamine and/or receptor), the compounds of the present invention may be used ~~for the manufacture of~~ pharmaceutical formulations for the treatment and prevention of diseases, damages and disorders, wherein biogenic amines and their receptors play an important role.

In view of the above explained favourable biological properties of the compounds of the present invention administration of the therapeutically effective amount of a compound of formula I provides an effective method of treatment of CNS diseases and disorders associated with fewer side effects due to their improved selectivity towards ~~the  $\sigma_{1\sigma 1}$~~  receptor and ~~the  $5-HT_{2A}$~~  and  $5-HT_{2C}$  serotonin receptors.

#### **Pharmaceutical Compositions**

In general, the compounds of the present invention may be used ~~for the manufacture of~~ pharmaceutical formulations that are used as antidepressants, anxiolytics, antipsychotics or as drugs for treating migraine.

Further, the compounds of the present invention may be used ~~for the manufacture of~~ pharmaceutical formulations for the treatment and prevention of diseases and disorders which are the result of disorders of neurochemical equilibrium in the central nervous system such as e.g. depression and modest depression, anxiety, bipolar disorders, sleeping disorders, sexual disorders, psychoses, borderline psychoses, schizophrenia, migraine, personality disorders and obsessive-compulsive disorders, social phobias or panic attacks, organic mental disorders in children, aggression, memory disorders and personality disorders in elderly people, addiction, obesity, bulimia and similar disorders, snoring, premenstrual troubles.

The present invention more specifically relates to an effective dose of the compounds which bind to serotonin, sigma, adrenergic, dopamine or muscarinic receptors and/or act as inhibitors of reabsorption of one or more biogenic amines (serotonin, dopamine, norepinephrine).

The pharmaceutical formulations are obtained by blending a therapeutically active amount of a certain substance as the active ingredient with a pharmaceutically acceptable carrier, which may have different forms depending on the desired administration route. These pharmaceutical formulations especially relate to oral, sublingual, rectal, percutaneous or parenteral administration route.

{W:\03818\0204416us0\00747961.DOC { }

The term "carrier" applied to pharmaceutical compositions of the invention refers to a diluent, excipient, or vehicle with which an active compound is administered. Such pharmaceutical carriers can be sterile liquids, such as water, saline solutions, aqueous dextrose solutions, aqueous glycerol solutions, and oils, including those of petroleum, animal, vegetable or synthetic origin, such as peanut oil, soybean oil, mineral oil, sesame oil and the like. However, since memantine is highly soluble, aqueous solutions are preferred. Suitable pharmaceutical carriers are described in "Remington's Pharmaceutical Sciences" by E.W. Martin, 18th Edition. Particularly preferred for the present invention are carriers suitable for immediate-release, i.e., release of most or all of the active ingredient over a short period of time, such as 60 minutes or less, and make rapid absorption of the drug possible.

A "pharmaceutically acceptable excipient" means an excipient that is useful in preparing a pharmaceutical composition that is generally safe, non-toxic and neither biologically nor otherwise undesirable, and includes an excipient that is acceptable for veterinary use as well as human pharmaceutical use. A "pharmaceutically acceptable excipient" as used in the present application includes both one and more than one such excipient.

"Treating" or "treatment" of a state, disorder or condition includes:

- (1) preventing or delaying the appearance of clinical symptoms of the state, disorder or condition developing in a mammal that may be afflicted with or predisposed to the state, disorder or condition but does not yet experience or display clinical or subclinical symptoms of the state, disorder or condition,
- (2) inhibiting the state, disorder or condition, i.e., arresting or reducing the development of the disease or at least one clinical or subclinical symptom thereof, or
- (3) relieving the disease, i.e., causing regression of the state, disorder or condition or at least one of its clinical or subclinical symptoms.

The benefit to a subject to be treated is either statistically significant or at least perceptible to the patient or to the physician.

A "therapeutically effective amount" means the amount of a compound that, when administered to a mammal for treating a state, disorder or condition, is sufficient to effect such treatment. The "therapeutically effective amount" will vary depending on the compound, the disease and its severity and the age, weight, physical condition and responsiveness of the mammal to be treated.

Dosages and administration regimen can be adjusted depending on the age, sex, physical condition as well as the benefit achieved by applying the compounds of the present invention and the side effects in the patient or the mammalian subject to be treated and the judgement of the physician, as is appreciated by those skilled in the art.

The term host or subject in need thereof as used herein refers to a mammal preferably a human.

#### **Biological Assays**

The effect of the compounds of the present invention on the neurochemical steady state was determined by *in vitro* investigations such as a radionuclide-marked radioligand binding assay for 5-HT<sub>2A</sub> (Bonhaus D. W. Br. J. Pharmacol. **1995**, 115:622; Saucier C. J. Neurochem. **1997**, 68:1998)

{W:\03818\0204416us0\00747961.DOC 19990901 11:00:00 AM 11/01/99 11:00:00 AM 11/01/99 11:00:00 AM }

and 5-HT<sub>2C</sub> receptors (Wolf W. A. *J. Neurochem.* **1997**, 69:1449), *in vitro* binding assay for the  $\sigma_1$  receptor (Thomson W. and Donn R. *Arthritis Res.* **2002**, 4: 302-306) and by *in vivo* investigations in a tail suspension test (Vogel H. G. and Vogel W. H. *Drug Discovery and Evaluation Pharmacological Assays*, Springer **1997**, 304), in amphetamine-induced hyperlocomotion in mice (Millan M.J. et al, **1998** *J Pharmacol. Exp. Ther.* 287: 167-186), in a forced swim test in mice (Porsolt R. D. et al. *Arch. Int. Pharmacodyn.* **1977**, 229:327–336), in meta-chlorophenyl piperazine (m-CPP) test on rats (*Drug Dev. Res.* **1989**, 18:119–144), and in apomorphine, tryptamine, norepinephrine (ATN) test in rats (*Arch. Int. Pharmacodyn.* **1977**, 227:238–253).

#### ***In vitro* method for determining affinity for binding to 5-HT<sub>2A</sub> and 5-HT<sub>2C</sub> receptors**

A small concentration of a radioligand having a great affinity for binding to a receptor was incubated with a tissue sample enriched with a certain receptor (1–5 mg of tissue) in a buffered medium (0.2–5 mL). Recombinant human HT<sub>2A</sub> and HT<sub>2C</sub> receptors were expressed in CHO-K1 or COS-7 cells and were also used for competitive binding. During incubation the radioligand bound to the receptor. When a binding balance was achieved, the receptors to which the radioligand was bound were separated from those to which said ligand was not bound, and the radioactivity of the receptor/radioligand complex was measured. The interaction of the tested compounds with receptors was tested in competitive binding experiments. Various concentrations of tested compounds were added to the incubation mixture containing a prepared tissue enriched with corresponding receptors and the radioligand. The radioligand binding was inhibited by the test compounds proportionally to the affinity of a certain compound for the receptor and to the concentration of the compound.

The radioligand used for the determination of binding to 5-HT<sub>2A</sub> receptor was [<sup>3</sup>H]-ketanserin and the tissue used was human cortex or recombinant 5-HT<sub>2A</sub> receptor expressed in CHO-K1 cells.

The radioligand used for the determination of binding to 5-HT<sub>2C</sub> receptor was [<sup>3</sup>H]-mesulergine and the tissue used was choroid plexus or recombinant 5-HT<sub>2C</sub> receptor expressed in CHO-K1 cells.

Compounds showing IC<sub>50</sub> and K<sub>i</sub> ~~in concentrations~~values lower than 1 μM, were considered to be active.

Compounds: 3-methyl-2-oxa-8-thia-1-aza-dibenzo[e,h]azulene and dimethyl-[3-(2-oxa-8-thia-1-aza-dibenzo[e,h]azulen-3-ylmethoxy)-propyl]-amine showed binding affinity to 5-HT<sub>2A</sub> and 5-HT<sub>2C</sub> serotonin receptors expressed as IC<sub>50</sub> value less than 200 nM and K<sub>i</sub> value less than 100 nM.

{W:\03818\0204416us0\00747961.DOC {FILED:\03818\0204416us0\00747961.DOC} }

Jurkat cell were grown in medium, RPMI supplemented with 10% fetal bovine serum, 100U/ml penicillin and 100µg/ml streptomycin, collected and their suspension homogenized. After centrifugation, membrane fraction was separated, resuspended in phosphate buffer (pH=7.5) and stored in small aliquots in liquid nitrogen until use.

Membranes were incubated with ligands in phosphate buffer for 3 hours at room temperature. After filter had been washed, radioactivity associated with the filter was determined by liquid scintillation spectrometry.

It is anticipated that similar results will be observed for other compounds of the invention.

Male CD1 mice of the weight of 20–25 g were used for the experiment. Groups of 10 animals were treated with the test compounds, imipramine (positive control) or the vehicle (negative control) by *per os* by gavage 30 min prior to testing to determine efficacy. On the day of the experiment the animals were placed into a glass cylinder (height 18.2 cm, diameter 13.3 cm) filled with water warmed to 22 °C to the height of 10 cm. The immobility defined as the end of the struggling of the animal and the beginning of floating, wherein the movements were reduced to those indispensable for the animal to keep its head over the water surface, started to be recorded after two minutes and then it was monitored during 4 minutes.

Male Balb/cJ mice of the weight of 20–25 g were used for the experiment. Groups of 9 animals were treated with the test compounds, imipramine (positive control) or the vehicle (negative control) by intraperitoneal injection, subcutaneous injection or per oral by gavage 30 min prior to testing to measure potential antidepressant activity. Mice were suspended from their tails at a height of about 90 cm and were observed for 5 minutes. The mice hanging fully motionless for 1 minute during the observation period were defined as depressive. In animals treated with a substance having an antidepressive action the period of immobility was shortened.

The percentage of animals showing a passive behaviour was calculated and compared with a control group treated with a vehicle. Significance of results was analysed using Fischer's exact test. The compounds that in a dose of 10 mg/kg reduced the immobility of animals for 40 % and more over a control group were considered to be active.

It is anticipated that similar results will be observed for other compounds of the invention.

Male Swiss OFA mice of a weight 30-35g were treated with either vehicle (saline) or test compounds 30 minutes prior to hyperlocomotion induction. Dexamphetamine sulphate was administered intraperitoneally at 2mg/kg. Thirty minutes later, animals were placed in a wooden box 80 x80 cm in a room with low light intensity (100 lux) for locomotor activity recording. Locomotor activity was determined during a 30 min period using a video image analyzer. Total duration of movement, occurrence of movement and total distance travelled were measured. Haloperidol was tested at the dose of 0,25 mg/kg (prepared in 0,5% methylcellulose) and served as reference substance.

Compounds were considered as active if in a dose of 10 mg/kg reduced amphetamine-induced hyperlocomotion in experimental animals for 30% and more when compared to vehicle treated control group.

It is anticipated that similar results will be observed for other compounds of the invention.







To a solution of 3-methyl-3,3a-dihydro-11-chloro-2-oxa-8-thia-1-aza-dibenzo[*e,h*]azulen-3-ol (**2a**) (0.08 mmole) in THF (5 mL), concentrated sulfuric acid (114  $\mu$ L) was added. The reaction mixture was stirred and heated under reflux for 5 hours, then it was cooled and the solvent was evaporated, water was added thereto and it was extracted with dichloromethane. The combined organic extracts were dried over anhydrous Na<sub>2</sub>SO<sub>4</sub> and evaporated under reduced pressure. After purification by chromatography on a silica gel column, an oily product was isolated;  
MS (*m/z*): 300.78 [MH<sup>+</sup>].

To a solution of 11H-dibenzo[b,f]oxepin-10-one oxime (1.91 mmole) in dry THF (10 mL) cooled to  $-78^{\circ}\text{C}$ , n-butyl lithium (4.10 mmole) was slowly added drop by drop. The reaction mixture was stirred for 15 minutes at this temperature, whereupon it was heated to  $0^{\circ}\text{C}$  and ethyl acetate (4.10 mmole) was added. The stirring of the reaction mixture was continued for one more hour at room temperature, whereupon water was added thereto and it was extracted with ethyl acetate. The combined organic extracts were dried over anhydrous  $\text{Na}_2\text{SO}_4$  and evaporated under reduced pressure. After purification by chromatography on a silica gel column, a crystalline product was isolated; MS ( $m/z$ ): 290.3  $[\text{MNa}^+]$ .

To a solution of 3-methyl-3,3a-dihydro-2,8-dioxo-1-aza-dibenzo[*e,h*]azulen-3-ol (**3a**) (0.1 mmole) in THF (7 mL), concentrated sulfuric acid (143  $\mu$ L) was added. The reaction mixture was stirred and heated under the reflux for 5 hours, then it was cooled and the solvent was evaporated, water was added thereto and it was extracted with dichloromethane. The combined organic extracts were dried over anhydrous Na<sub>2</sub>SO<sub>4</sub> and evaporated under reduced pressure. After purification by chromatography on a silica gel column, an oily product was isolated;  
MS (*m/z*): 250.27 [MH<sup>+</sup>].

{W:\03818\0204416us0\00747961.DOC 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60 }

*Dimethyl-[3-(2-oxa-8-thia-1-aza-dibenzo[e,h]azulen-3-ylmethoxy)-propyl]-amine (7)*

To a solution of 3-dimethylaminopropylchloride-hydrochloride (2.16 mmole) in 50 % sodium hydroxide (1.9 mL), a catalytic amount of benzyltriethylammonium chloride and a solution of 1-bromomethyl-2-oxa-8-thia-1-aza-dibenzo[*e,h*]azulene (**4**) (0.15 mmole) in toluene (10 mL) were added. The reaction mixture was heated under vigorous stirring and reflux for 3 hours, then it was cooled to room temperature, diluted with water and extracted with dichloromethane. The organic extract was washed with water, dried over anhydrous Na<sub>2</sub>SO<sub>4</sub> and evaporated under reduced pressure. After purification by chromatography on a silica gel column, a crystalline product was isolated; MS (*m/z*): 367.2 [MH<sup>+</sup>].

### Example 8

*Dimethyl-[2-(2-oxa-8-thia-1-aza-dibenzo[e,h]azulen-3-ylmethoxy)-ethyl]-amine (8)*

According to the process described in Example 7, starting from 1-bromomethyl-2-oxa-8-thia-1-azadibenzo[*e,h*]azulene (**4**) (0.20 mmole) and 2-dimethylamino-ethylchloride-hydrochloride (2.85 mmole), an oily product was obtained;

<sup>1</sup>H NMR (ppm, CDCl<sub>3</sub>): 2.39 (s, 6H), 2.69-2.72 (t, 2H), 3.83-3.87 (t, 2H), 4.79 (s, 2H), 7.35-7.89 (m, 8H);

MS (*m/z*): 353.2 [MH<sup>+</sup>], 375.2 [MNa<sup>+</sup>].

### Example 9

*Dimethyl-[2-(11-chloro-2-oxa-8-thia-1-aza-dibenzo[e,h]azulen-3-ylmethoxy)-ethyl]-amine (9)*

According to the process described in Example 7, starting from 1-bromomethyl-11-chloro-2-oxa-8-thia-1-aza-dibenzo[*e,h*]azulene (**5**) (0.18 mmole) and 2-dimethylaminoethylchloride-hydrochloride (2.56 mmole), an oily product was obtained;

MS (*m/z*): 387.65 [MH<sup>+</sup>].

### Example 10

*Dimethyl-[3-(11-chloro-2-oxa-8-thia-1-aza-dibenzo[e,h]azulen-3-ylmethoxy)-propyl]-amine (10)*

According to the process described in Example 7, starting from 1-bromomethyl-11-chloro-2-oxa-8-thia-1-aza-dibenzo[*e,h*]azulene (**5**) (0.18 mmole) and 2-dimethylaminopropylchloride-hydrochloride (2.56 mmole), an oily product was obtained;

MS ( $m/z$ ): 401.65 [MH<sup>+</sup>].

### Example 11

*Dimethyl-[2-(2,8-dioxa-1-aza-dibenzo[e,h]azulen-3-ylmethoxy)-ethyl]-amine (11)*

*Dimethyl-[3-(2,8-dioxa-1-aza-dibenzo[e,h]azulen-3-ylmethoxy)-propyl]-amine (12)*

According to the process described in Example 7, starting from 1-bromomethyl-2,8-dioxo-1-azadibenzo[*e,h*]azulene (6) (0.25 mmole) and 2-dimethylamino-propylchloride-hydrochloride (3.42 mmole), an oily product was obtained;  
MS (*m/z*): 351.2 [MH<sup>+</sup>].

*11H-dibenzo[b,f]thiepin-10-one oxime*

11H-dibenzo[*b,f*]thiepin-10-one (J.O. Jilek et al. *Mh. Chem.* **96** (1965) 182-207) (2.21 mmole) was dissolved in absolute ethanol (4.26 mL) and water (1.28 mL) under stirring and gentle heating. To the solution of ketone, aminehydroxide hydrochloride (4.42 mmole) and sodium acetate (4.42 mmole) were added. The reaction mixture was stirred and heated under reflux for 2 hours. After the completion of the reaction, 30 % ethanol (2 mL) was added into the hot reaction mixture and it was left to cool to room temperature. If no precipitation occurred, the solvent was evaporated under reduced pressure and the residue after evaporation was dissolved in water, extracted with dichloromethane, dried over anhydrous Na<sub>2</sub>SO<sub>4</sub> and evaporated under reduced pressure. After purification by chromatography on a silica gel column, a crystalline product was isolated;

<sup>1</sup>H NMR (ppm, CDCl<sub>3</sub>): 3.65 (bs, 1H), 4.34 (s, 2H), 7.18-8.06 (m, 8H);

MS (*m/z*): 242.0 [MH<sup>+</sup>], 264.0 [MNa<sup>+</sup>], 296.0 [MNa<sup>+</sup> + MeOH].

*8-chloro-11H-dibenzo[b,f]thiepin-10-one oxime*

11-chloro-11H-dibenzo[*a,f*]thiepin-10-one (J.O. Jilek et al. *Mh. Chem.* **96** (1965) 182-207) (1,47 mmole) was dissolved in absolute ethanol (2.84 mL) and water (0.9 mL) under stirring and gentle heating. To the solution of ketone, aminehydroxide hydrochloride (2.95 mmole) and sodium acetate (2.95 mmole) were added. The reaction mixture was stirred and heated under reflux for 2 hours. After the completion of the reaction, 30 % ethanol (1 mL) was added into the hot reaction mixture and it was left to cool to room temperature. If no precipitation occurred, the solvent was evaporated under

11H-dibenzo[*b,f*]oxepin-10-one (I. Ueda et al. *Chem. Pharm. Bull.* **23** (10) 2223-2231 (1975)) (4.42 mmole) was dissolved in absolute ethanol (8.52 mL) and water (2.56 mL) under stirring and gentle heating. To the solution of ketone, aminehydroxide hydrochloride (8.84 mmole) and sodium acetate (8.84 mmole) were added. The reaction mixture was stirred and heated under reflux for 2 hours. After the completion of the reaction, 30 % ethanol (4 mL) was added into the hot reaction mixture and it was left to cool to room temperature. If no precipitation occurred, the solvent was evaporated under reduced pressure and the residue after evaporation was dissolved in water, extracted with dichloromethane, dried over anhydrous Na<sub>2</sub>SO<sub>4</sub> and evaporated under reduced pressure. After purification by chromatography on a silica gel column, a crystalline product was isolated; MS (*m/z*): 226.0 [MH<sup>+</sup>].

Compound	Structure	Name
<b>1a</b>		3-Methyl-3,3a-dihydro-2-oxa-8-thia-1-aza-dibenzo[e,h]azulen-3-ol
<b>2a</b>		11-Chloro-3-methyl-3,3a-dihydro-2-oxa-8-thia-1-aza-dibenzo[e,h]azulen-3-ol
<b>3a</b>		3-Methyl-3,3a-dihydro-2,8-dioxo-1-aza-dibenzo[e,h]azulen-3-ol
<b>1</b>		3-Methyl-2-oxa-8-thia-1-aza-dibenzo[e,h]azulene

2		11-Chloro-3-methyl-2-oxa-8-thia-1-aza-dibenzo[e,h]azulene
3		3-Methyl-2,8-dioxa-1-aza-dibenzo[e,h]azulene
4		3-Bromomethyl-2-oxa-8-thia-1-aza-dibenzo[e,h]azulene
5		3-Bromomethyl-2,8-dioxa-1-aza-dibenzo[e,h]azulene
6		3-Bromomethyl-11-chloro-2-oxa-8-thia-1-aza-dibenzo[e,h]azulene
7		Dimethyl-[3-(2-oxa-8-thia-1-aza-dibenzo[e,h]azulen-3-ylmethoxy)-propyl]-amine
8		Dimethyl-[2-(2-oxa-8-thia-1-aza-dibenzo[e,h]azulen-3-ylmethoxy)-ethyl]-amine
9		2-[(11-Chloro-2-oxa-8-thia-1-aza-dibenzo[e,h]azulen-3-ylmethoxy)-ethyl]-dimethyl-amine
10		3-[(11-Chloro-2-oxa-8-thia-1-aza-dibenzo[e,h]azulen-3-ylmethoxy)-propyl]-dimethyl-amine
11		2-(2,8-Dioxa-1-aza-dibenzo[e,h]azulen-3-ylmethoxy)-ethyl-dimethyl-amine
12		3-(2,8-Dioxa-1-aza-dibenzo[e,h]azulen-3-ylmethoxy)-propyl-dimethyl-amine